arXiv:math-ph/0703066v1 21 Mar 2007

Multisoliton solutions of 3,4,6 waves problems connectet with semisimlealgebras of the second rank $A_3, B_2 = C_2, G_2$

A. N. Leznov*

Abstract

With each semisimple algebra it is possible to connect the system of interacting waves. The number of interacting fields coincides with the number of positive roots of corresponding semisimple algebra. Multisoliton solution of such kind problem are represented in explicit form for the algebras of second rank.

1 Introduction

In the present paper we would like to constract multisoliton solutions for n-waves interacting system in explicit form, using technique of discrete transformation theory introduced in [1]. The equation described n- waves interacting system are the following [1]

$$[(dh), f_t] + [(ch), f_x] = [[(dh), f], [(ch), f]] \tag{1}$$

where f is algebra-valued unknown function (taking values in arbitrary semisimple algebra), (x, t) independent arguments of the problem. In component form (1) looks as

$$(c_R \frac{\partial}{\partial t} + d_R \frac{\partial}{\partial x}) f_R = \sum_P (c_{R-P} d_P - d_{R-P} c_P) f_{R-P} f_P \tag{2}$$

where by indexes P, R are defined the set of all roots of semisimple algebra c_P, d_P values of cartan elements $(ch), (dh)$ on these roots. The case of A_2 algebra was considered in details in [2] and multisoliton solutions in determinant form for this problem was present in [3]. In [4] this results was rediscovered in the form which allow generalisation on the case of all semisimple algebras of the second rank (see sections below).

---

*Universidad Autonoma del Estado de Morelos, CCICAp,Cuernavaca, Mexico

2 Solution of main equations in the case of of lower triangular algebra

In this section we find solution of (2) under additional condition $f^+_R=0$ for all roots of semisimple algebra.

2.0.1 Simple roots

For each simple root equations (2) looks as

$$(c_\alpha \frac{\partial}{\partial t}+d_\alpha \frac{\partial}{\partial x})f^-_\alpha=0$$

with the obvious solution $f^-_\alpha=\Phi_\alpha(d_\alpha t-c_\alpha x)$. In the problem of the present paper for us it be essential parametrization of f^-_α in the form (of Lapalce transformation)

$$f^-_\alpha=\int d\lambda_\alpha s^\alpha(\lambda_\alpha)e^{\lambda_\alpha(d_\alpha t-c_\alpha x)} \qquad (3)$$

Thus arbitarary simple root α of algebra is connected with arbitrary function of one variable $s^\alpha(\lambda_\alpha)$.

2.0.2 Complicate root constructed from two simple ones

The equation for such component is the following

$$((c_\alpha+c_\beta)\frac{\partial}{\partial t}+(d_\alpha+d_\beta)\frac{\partial}{\partial x})f^-_{\alpha,\beta}=(c_\alpha d_\beta-d_\alpha c_\beta)f^-_\alpha f^-_\beta$$

Without any difficulties it is possible verified that solution of this equation is the following

$$f^-_{\alpha,\beta}=\int d\lambda_\alpha d\lambda_\beta \frac{s^\alpha(\lambda_\alpha)s^\beta(\lambda_\beta)}{\lambda_\alpha-\lambda_\beta}e^{t(d_\alpha\lambda_\alpha+d_\beta\lambda_\beta)-x(c_\alpha\lambda_\alpha+c_\beta\lambda_\beta)} \qquad (4)$$

Plus solution of homogineous equation.

2.0.3 Complicate root constructed from three simple ones

It is possible two cases, when the fird root γ not equal to one of the previous ones α or β or it coincedes with one of them. At first let us consider the first possibility. The equation (2) in this case looks as

$$((c_\alpha+c_\beta+c_\gamma)\frac{\partial}{\partial t}+(d_\alpha+d_\beta+d_\gamma)\frac{\partial}{\partial x})f^-_{\alpha,\beta,\gamma}=$$

$$(c_\alpha(d_\beta+d_\gamma)-d_\alpha(c_\beta+c_\gamma))f^-_\alpha f^-_{\beta,\gamma}+((c_\alpha+c_\beta)d_\gamma)-(d_\alpha+d_\beta)c_\gamma)f^-_{\alpha,\beta}f^-_\gamma \qquad (5)$$

By direct check we find solution of the last equation in the form

$$f^-_{\alpha,\beta,\gamma}=\int d\lambda_\alpha d\lambda_\beta d\lambda_\gamma \frac{s^\alpha(\lambda_\alpha)s^\beta(\lambda_\beta)s^\gamma(\lambda_\gamma)}{(\lambda_\alpha-\lambda_\beta)(\lambda_\beta-\lambda_\gamma)}e^{t(d_\alpha\lambda_\alpha+d_\beta\lambda_\beta+d_\gamma\lambda_\gamma)-x(c_\alpha\lambda_\alpha+c_\beta\lambda_\beta+c_\gamma\lambda_\gamma)}$$

In the second case, when for instance $\beta=\gamma$ the defining equation (5)does not contain one of the terms and in this case solution looks as

$$f^-_{\alpha,\beta,\beta}=\int d\lambda_\alpha d\lambda_{\beta_1} d\lambda_{\beta_2} \frac{s^\alpha(\lambda_\alpha)s^\beta(\lambda_{\beta_1})s^\beta(\lambda_{\beta_2})}{(\lambda_\alpha-\lambda_{\beta_1})(\lambda_\alpha-\lambda_{\beta_2})}e^{t(d_\alpha\lambda_\alpha+d_\beta(\lambda_{\beta_1}+\lambda_{\beta_2}))-x(c_\alpha\lambda_\alpha+c_\beta(\lambda_{\beta_1}+\lambda_{\beta_2}))}$$

3 The case of A_2 algebra

This section is written only for convineous of the reader and contain necessary for what follows results of [2],[4]. The results conserninining B_2 and G_2 algebras will be presented below in the same form.

Algebra A_2 has the following Cartan matrix and basic commutation relations between two generators of the simple roots $X^{\pm}_{1,2}$ and its Cartan elements $h_{1,2}$

$$k=\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix},\quad \begin{pmatrix} [h_1,X^{\pm}_1]=\pm 2X^{\pm}_1 & [h_1,X^{\pm}_2]=\mp X^{\pm}_2 \\ [h_2,X^{\pm}_1]=\mp X^{\pm}_2 & [h_2,X^{\pm}_2]=\pm 2X^{\pm}_2 \end{pmatrix}$$

Arbitrary element of the algebra may be represented as (up to Cartan elements which are absent in the problem under consideration)

$$f=f^+_{1.1}X^+_{\alpha_1+\alpha_2}+f^+_{0.1}2X^+_{\alpha_2}+f^+_{1.0}X^+_{\alpha_1}+f^-_{1.0}X^-_{\alpha_1}+f^-_{0.1}X^-_{\alpha_2}+f^-_{1.1}X^-_{\alpha_1+\alpha_2}$$

$\alpha_{1,2}$ are the indexes of si mple roots. $X^+_{\alpha_1+\alpha_2}\equiv[X^+_2,X^+_1]$.

In these notations the system of equations (2) looks as

$$\begin{aligned} D_{1,0}f^+_{1.0}&=f^+_{1.1}f^-_{0.1}, \quad D_{1,0}f^-_{1.0}=f^-_{1.1}f^+_{0.1} \\ D_{0,1}f^+_{0.1}&=f^+_{1.1}f^-_{1.0}, \quad D_{0,1}f^-_{0.1}=f^-_{1.1}f^+_{1.0} \\ D_{1,1}f^+_{1.1}&=-f^+_{0.1}f^+_{1.0}, \quad D_{1,1}f^-_{1.1}=-f^-_{0.1}f^-_{1.0} \end{aligned} \tag{6}$$

where operators of differenciation are the following ones $D_{i,j}=\frac{(ic_1+jc_2)}{\delta}\frac{\partial}{\partial t}+\frac{(id_1+jd_2)}{\delta}\frac{\partial}{\partial x}$, $\delta\equiv(c_1d_2-c_2d_1)$.

The discrete transformation of this system are the following ones [2]

3.0.4 T_3

The system (6) is invariant with respect to the following transformation T_3

$$\tilde{f}^+_{1.1}=\frac{1}{f^-_{1.1}},\quad \tilde{f}^+_{1.0}=-\frac{f^-_{0.1}}{f^-_{1.1}},\quad \tilde{f}^+_{0.1}=\frac{f^-_{1.0}}{f^-_{1.1}}$$

$$\tilde{f}^-_{0.1} = -f^-_{1.1} D_{1.0} \frac{f^-_{0,1}}{f^-_{1.1}}, \quad \tilde{f}^-_{1.0} = f^-_{1.1} D_{0,1} \frac{f^-_{1.0}}{f^-_{1.1}} \tag{7}$$

$$\frac{\tilde{f}^-_{1.1}}{f^-_{1.1}} = f^+_{1.1} f^-_{1.1} - D_{1,0} D_{0,1} \ln f^-_{1.1}$$

3.0.5 T_2

The system (6) is invariant with respect to the following transformation T_2

$$\tilde{f}^+_{0.1} = \frac{1}{f^-_{0.1}}, \quad \tilde{f}^-_{1.0} = -\frac{f^-_{1.1}}{f^-_{0.1}}, \quad \tilde{f}^+_{12} = \frac{f^+_{1.0}}{f^-_{0.1}}$$

$$\tilde{f}^+_{1.0} = -f^-_{0.1} D_{1,1} \frac{f^+_{1.0}}{f^-_{0.1}}, \quad \tilde{f}^-_{1.1} = -f^-_{0.1} D_{1,0} \frac{f^-_{1.1}}{f^-_{0.1}} \tag{8}$$

$$\frac{\tilde{f}^-_{0.1}}{f^-_{0.1}} = f^+_{0.1} f^-_{0.1} + D_{1,0} D_{1,1} \ln f^-_{0.1}$$

3.0.6 T_1

The system (6) is invariant with respect to the following transformation T_1

$$\tilde{f}^+_{1.0} = \frac{1}{f^-_{1.0}}, \quad \tilde{f}^-_{0.1} = \frac{f^-_{1.1}}{f^-_{1.0}}, \quad \tilde{f}^+_{1.1} = -\frac{f^+_{0.1}}{f^-_{1.0}}$$

$$\tilde{f}^+_{0.1} = f^-_{0.1} D_{1,1} \frac{f^+_{0.1}}{f^-_{1.0}}, \quad \tilde{f}^-_{1.1} = f^-_{1.0} D_{0,1} \frac{f^-_{1.1}}{f^-_{1.0}} \tag{9}$$

$$\frac{\tilde{f}^-_{1.0}}{f^-_{1.0}} = f^+_{1.0} f^-_{1.0} + D_{0,1} D_{1,1} \ln f^-_{1.0}$$

3.0.7 General properties of discrete transformations

Three above transformations are invertable. This means f may be expressed algebraicaly in terms of $\tilde{f}$. Exept of this $T_3 = T_1 T_2 = T_2 T_1$, what means that all discrete transformation are mutually commutative. This in its turn means that arbitrary discrete transformation may be represented in a form $T = T_1^{n_1} T_2^{n_2}$ [2].

3.0.8 Result of consequent application of some number of discrete transformations

In determinant form result was found in [2]. In [4] it was rediscovered in the form which can be generalised to the case of arbitrary semi-simple algebra of second rank (see sections below).

Let us rewrite (3) and (4) in a form

$$f^-_{1.0}=\int d\lambda p(\lambda)e^{\lambda(d_1t-c_1x)}\equiv\int d\lambda P(\lambda),\quad f^-_{0.1}=\int d\mu q(\mu)e^{\mu(d_2t-c_2x)}\equiv\int d\mu Q(\mu),$$

$$f^-_{1.1}=\int d\lambda\int d\mu\frac{p(\lambda)q(\mu)}{\lambda-\mu}e^{\lambda(d_1t-c_1x)+\mu(d_2t-c_2x)}\equiv\int d\lambda\int d\mu\frac{P(\lambda)Q(\mu)}{\lambda-\mu}$$

and choose initial condition in a form

$$f^+_{1.0}=f^+_{0.1}=f^+_{1.1}=0$$

To solution of this kind application of each inverse transformation T_i^{-1} is meanless wia zeros in denumerators.

Let us introduce determining function $U(n_1,n_2)$

$$U(n_1,n_2)=\int\prod_{i=1}^{n_1}P(\lambda_i)d\lambda_i\prod_{k=1}^{n_2}Q(\mu_k)d\mu_k\frac{W^2_{n_1}(\lambda)W^2_{n_2}(\mu)}{\prod_{i,k}(\lambda_i-\mu_k)}\tag{10}$$

where W_n is Vandermond determinant constructed from $n_{1,2}$-variables λ or μ.

Then result of appliucation n_1 times discrete transformation T_1 and n_2 times discrete transformation T_2 looks as [4]

$$\begin{aligned}f^+_{1.0}&=\frac{U(n_1-1,n_2)}{U(n_1,n_2)},\quad f^-_{1.0}=\frac{U(n_1+1,n_2)}{U(n_1,n_2)},\\ f^+_{0.1}&=\frac{U(n_1,n_2-1)}{U(n_1,n_2)},\quad f^-_{0.1}=\frac{U(n_1,n_2+1)}{U(n_1,n_2)},\\ f^+_{1.1}&=\frac{U(n_1-1,n_2-1)}{U(n_1,n_2)},\quad f^-_{1.1}=\frac{U(n_1+1,n_2+1)}{U(n_1,n_2)}\end{aligned}\tag{11}$$

In the case of integrable systems connected with A_1 algebra Vandermond form for soliton solutions was introduced and used in [6].

Let us assume that initial functions has the form $P(\lambda)=\sum_{i=1}^{n_1}\delta(\lambda-l_k)e^{l_k(d_1t-c_1x)}$ and $Q(\mu)=\sum_{i=1}^{n_2}\delta(\mu-m_i)e^{m_i(d_2t-c_2x)}$, where $\delta(x)$ is the ussual Dirac δ function. In this case as it follows from (11) on the n_1,n_2 step of discrete transformation $f^-_{1.0}=f^-_{0.1}=f^-_{1.1}=0$ the chain is interrupted on the second end. And from results of [2] conditions of reality lead directly to soliton solutions.

4 The case of B_2 algebra

Algebra B_2 (eqiuvalent to the algebra of the group of 5-dimentional rotations) has the following Cartan matrix and basics comutation relations

$$k=\begin{pmatrix}2&-2\\-1&2\end{pmatrix},\quad\begin{pmatrix}[h_1,X^\pm_1]=\pm2X^\pm_1&[h_1,X^\pm_2]=\mp X^\pm_2\\ [h_2,X^\pm_1]=\mp2X^\pm_2&[h_2,X^\pm_2]=\pm2X^\pm_2\end{pmatrix}$$

Arbitrary element of this algebra can be parametrised as (up to elements taking values in Cartan subalgebra)

$$f = f^+_{1.2} X^+_{\alpha_1+2\alpha_2} + f^+_{1,1} X^+_{\alpha_1+\alpha_2} + f^+_{0.1} X^+_{\alpha_2} + f^+_{1.0} X^+_{\alpha_1} + f^-_{1.0} X^-_{\alpha_1} + f^-_{0.1} X^-_{\alpha_2} + f^-_{1.1} X^-_{\alpha_1+\alpha_2} + f^-_{1.2} X^-_{\alpha_1+2\alpha_2}$$

$\alpha_{1,2}$ are the indexes of simple roots. In these notations the system of equations (2) looks as

$$\begin{gathered}
D_{1,0} f^+_{1.0} = 2 f^+_{1.1} f^-_{0.1}, \quad D_{1,0} f^-_{1.0} = 2 f^-_{1.1} f^+_{0.1} \\
D_{0,1} f^+_{0.1} = f^+_{1.1} f^-_{1.0} + f^+_{1.2} f^-_{1.1}, \quad D_{0,1} f^-_{0.1} = f^-_{1.1} f^+_{1.0} + f^-_{1.2} f^+_{1.1} \\
D_{1,1} f^+_{1.1} = -f^+_{0.1} f^+_{1.0} + f^+_{1.2} f^-_{0.1}, \quad D_{1,1} f^-_{1.1} = -f^-_{0.1} f^-_{1.0} + f^-_{1.2} f^+_{0.1} \\
D_{1,2} f^+_{1,2} = -2 f^+_{1.1} f^+_{0.1}, \quad D_{1,2} f^-_{1.2} = -2 f^-_{1.1} f^-_{0.1}
\end{gathered} \tag{12}$$

where $D_{i,j} = (\frac{ic_1+jc_2}{\delta} \frac{\partial}{\partial t} + \frac{id_1+jd_2}{\delta} \frac{\partial}{\partial x})$, $\delta \equiv (c_1 d_2 - c_2 d_1)$. The above system of equations (12) is the wide system of interaction of eight fields $f^\pm_{1.0}, f^\pm_{0.1}, f^\pm_{1.1}, f^\pm_{1.2}$. This system is obviously allow reducing $(f^-)^H = f^+$ (under additional assumtion that all operators of differentiations are real). Exactly such kind solutions of this system is intersting for applications and will be the point of investigation in the present section.

The discrete transformation to the such kind of the systems in the case of arbitrary semisimple algebra was found in [1]. The formulae below are partial case of application general construction for the case of B_2 algebra. They can be easely checked by direct not combersome calculation.

4.1 Discrete transformation T_M

The system (12) invariant with respect to following transformation T_M (discrete transformation of the maximal root of B_2 algebra)

$$\tilde{f}^+_{1.2} = \frac{1}{f^-_{1.2}}, \quad \tilde{f}^+_{0.1} = \frac{f^-_{1.1}}{f^-_{1.2}}, \quad \tilde{f}^+_{1.1} = -\frac{f^-_{0.1}}{f^-_{1.2}}, \quad \tilde{f}^+_{1.0} = f^+_{1.0} + \frac{(f^-_{0.1})^2}{f^-_{1.2}}, \quad \tilde{f}^-_{1.0} = f^-_{1.0} - \frac{(f^-_{1.1})^2}{f^-_{1.2}}$$

$$\tilde{f}^-_2 = -D_{1,0} f^-_2 - f^+_{1.1} f^-_{1.2} + \frac{1}{2} f^-_{0.1} D_{1,0} \ln f^-_{1.2}, \quad \tilde{f}^-_{1.1} = -D_{1,0} f^-_{1.1} + f^+_{0.1} f^-_{1.2} + \frac{1}{2} f^-_{1.1} D_{1,0} \ln f^-_{1.2}$$

$$\frac{\tilde{f}^-_{1.2}}{f^-_{1.2}} = \frac{1}{4} D_{1,0} D_{1,0} \ln f^-_{1.2} + \frac{f^-_{1.1} D_{1,0} f^-_{0.1} - f^-_{0.1} D_{1,0} f^-_{1.1}}{2 f^-_{1.2}} + f^+_{1.2} f^-_{1.2} + f^+_{1.1} f^-_{1.1} + f^+_{0.1} f^-_{0.1}$$

By direct not combersome calculations it is possible verified that $\tilde{f}$ satisfy (12) if f is solution of this system.

4.2 Discrete transformation $T_{1.0}$

$T_{1.0}$-discrete transformation of the first (α_1) root of B_2 algebra

$$\tilde{f}^+_{1.0} = \frac{1}{f^-_{1.0}}, \quad \tilde{f}^-_{0.1} = \frac{f^-_{1.1}}{f^-_{1.0}}, \quad \tilde{f}^+_{1.1} = -\frac{f^+_{0.1}}{f^-_{1.0}}, \quad \tilde{f}^+_{1.2} = f^+_{1.2} + \frac{(f^+_{0.1})^2}{f^-_1}, \quad \tilde{f}^-_{1.2} = f^-_{1.2} - \frac{(f^-_{1.1})^2}{f^-_{1.0}}$$

$$\tilde{f}^+_{0.1} = D_{1,2} f^+_{0.1} - f^+_{1.1} f^-_{1.0} - \frac{1}{2} f^+_{0.1} D_{1,2} \ln f^-_{1.0}, \quad \tilde{f}^-_{1.1} = D_{1,2} f^-_{1,1} + f^-_{0.1} f^-_{1.0} - \frac{1}{2} f^-_{1.1} D_{1,2} \ln f^-_{1.0} \tag{13}$$

$$\frac{\tilde{f}^-_{1.0}}{f^-_{1.0}} = \frac{1}{4} D_{1,2} D_{1.2} \ln f^-_{1.0} + \frac{f^-_{1.1} D_{1,2} f^+_{0.1} - f^+_{0.1} D_{1,2} f^-_{1.1}}{2 f^-_{1.0}} + f^+_{1.0} f^-_{1.0} - f^+_{1.1} f^-_{1.1} + f^+_{0.1} f^-_{0.1}$$

The most important propertie of this transformation consists in the fact that if we begin from the initial solution $f^+_{1.0} = f^+_{0.1} = f^+_{1.1} = f^+_{1.2} = 0$ the last three function contains there zero values after discrete transformation $T_{1.0}$ as it follows from (13).

4.3 Inverse discrete transformation $T^{-1}_{1.0}$

To construct inverse $T^{-1}_{1.0}$ it is necessary formulae (13) resolve with respect to untilded functions. Result is the the following one

$$f^-_{1.0} = \frac{1}{\tilde{f}^+_{1.0}}, \quad f^+_{0.1} = -\frac{\tilde{f}^+_{1.1}}{\tilde{f}^+_{1.0}}, \quad f^-_{1.1} = \frac{\tilde{f}^-_{0.1}}{\tilde{f}^+_{1.0}}, \quad f^+_{1.2} = \tilde{f}^+_{1.2} - \frac{(\tilde{f}^+_{1.1})^2}{\tilde{f}^+_{1.0}}, \quad f^-_{1.2} = \tilde{f}^-_{1.2} + \frac{(\tilde{f}^-_{0.1})^2}{\tilde{f}^+_{1.0}}$$

$$f^-_{0.1} = -D_{1,2} \tilde{f}^-_{0.1} - \tilde{f}^-_{1.1} \tilde{f}^+_{1.0} - \frac{1}{2} \tilde{f}^-_{0.1} D_{1,2} \ln \tilde{f}^+_{1.0}, \quad f^+_{1.1} = -D_{1,2} \tilde{f}^+_{1.1} + \tilde{f}^+_{0.1} \tilde{f}^+_{1.0} - \frac{1}{2} \tilde{f}^+_{1.1} D_{1,2} \ln \tilde{f}^+_{1.0} \tag{14}$$

$$\frac{f^+_{1.0}}{\tilde{f}^+_{1.0}} = \frac{1}{4} D_{1,2} D_{1,2} \ln \tilde{f}^+_{1.0} - \frac{\tilde{f}^+_{1.1} D_{1,2} \tilde{f}^-_{0.1} - \tilde{f}^-_{0.1} D_{1,2} \tilde{f}^+_{1.1}}{2 \tilde{f}^+_{1.0}} + \tilde{f}^+_{1.0} \tilde{f}^-_{1.0} + \tilde{f}^+_{1.1} \tilde{f}^-_{1.1} + \tilde{f}^+_{0.1} \tilde{f}^-_{0.1}$$

4.4 Discrete transformation $T_{2\alpha_2}$

In the case of B_2 algebra it is not possible to coinside with the help of the Weil group the maximal root with $X^{\pm}_{\alpha_2}$. But having in mind T_M and T_1 it is possible to construct dicrete transformation $T_{2\alpha_2} \equiv T_{\alpha_1+2\alpha_2} T^{-1}_{\alpha_1} = T_M T^{-1}_1$ which we will call discrete transformation of the second root and which looks as

$$\tilde{f}^-_{1.0} = \frac{1}{f^+_{1.0} + \frac{(f^-_{0.1})^2}{f^-_{1.2}}}, \quad \tilde{f}^-_{1.1} = \frac{-D_{1.0} f^-_{0.1} - f^+_{1.1} f^-_{1.2} + \frac{1}{2} f^-_{0.1} D_{1.0} \ln f^-_{1.2}}{f^+_{1.0} + \frac{(f^-_{0.1})^2}{f^-_{1.2}}}, \quad \tilde{f}^+_{0.1} = \frac{f^-_{0.1}}{f^+_{1.0} f^-_{1.2} + (f^-_{0.1})^2}$$

$$\tilde{f}^+_{1.2} = \frac{f^+_{1.0}}{f^+_{1.0} f^-_{1.2} + (f^-_{0.1})^2}, \quad \tilde{f}^+_{1.1} = \frac{f^+_{1.0} D_{1.2} f^-_{0.1} - f^-_{1.1} (f^+_{1.0})^2 - \frac{1}{2} f^-_{0.1} D_{1.2} f^+_{1.0}}{f^+_{1.0} + \frac{(f^-_{0.1})^2}{f^-_{1.2}}}$$

From the last formulae it follows that $T_{2\alpha_2}$ conserve zero values for wave functions $f^+_{1.0} = f^+_{1.1} = f^+_{1.2} = 0$ under initial conditions under consideration. By this reason we present further formulae after substitution in them these values. Because the general formulae are suffitiently combersom.

$$\frac{\tilde{f}^-_{0.1}}{f^-_{0.1}} = D^2_{1,0} \ln f^-_{0.1} + f^-_{0.1} f^+_{0.1}, \quad \tilde{f}^+_{0.1} = \frac{1}{f^-_{0.1}}$$

This is the equations of the ussual Toda lattice solution of which are well known. And at last after some number of manipulations we obtain

$$\tilde{f}^-_{1.2} = \frac{1}{4} D^2_{1,0} f^-_{1.2} + \frac{f^-_{1.1} D_{1,0} f^-_{0.1} - f^-_{0.1} D_{1,0} f^-_{1.1}}{2} +$$
$$\frac{D_{1,0} f^-_{0.1} D_{1,0} f^-_{0.1}}{(f^-_{0.1})^2} f^-_{1.2} - \frac{D_{1,0} f^-_{0.1} D_{1.0} f^-_{1.2}}{f^-_{0.1}} + f^+_{0.1} f^-_{0.1} f^-_{1.2}$$

4.5 Linear integrable chain of the second simple root in the field of Toda lattice

Let us consider consequent action of discrete transformation of the previous subsection on the initial solution $f^+_{1.0} = f^+_{0.1} = f^+_{1.1} = f^+_{1.2} = 0$. As it fololows from explicit formulae for discrete transformation zero values for $f^+_{1.0} = f^+_{1.1} = f^+_{1.2} = 0$ are conserved. Equations for $f^-_{0.1}, f^+_{0.1}$ functions are typical one dimentional Toda chain with well known solution. Result of the application n-times of Toda discrete transformation to initial functon $f^-_{0.1} \equiv r$ is the following

$$f^+_{0.1} = \frac{Det_{n-1}}{Det_n}, \quad f^-_{0.1} = \frac{Det_{n+1}}{Det_n} \tag{15}$$

where Det_n are the the main minores from left upper corner of the matrix

$$\Delta = \begin{pmatrix} r & r_1 & r_{11} & ..... \\ r_1 & r_{11} & r_{111} & ..... \\ r_{11} & r_{111} & r_{1111} & ..... \\ ...... & .... & .... & ..... \end{pmatrix} \tag{16}$$

where $r_1 \equiv D_{1,0} r$ and so on. In these notations equations of Toda latice looks as

$$D^2_{1,0}(\ln D_n) = \frac{Det_{n-1} Det_{n+1}}{Det^2_n} \tag{17}$$

Initial function of the second simple root $r = f^-_{0.1} = \int d\mu q(\mu) e^{\mu(d_2 t - c_2 x)} \equiv \int d\mu Q$. In Q all dependence on arguments x, t is included. Obviously $D_{0,1} Q = 0$. In these notations Det_n may be presented in the form of multidimensional integrale

$$Det_n = \frac{1}{n!} \int \prod_{i=1}^{n} d\mu_i Q_{\mu_i} W^2_n(\mu_1, .., \mu_n)$$

where $W^2_n(\mu_1, .., \mu_n)$ is squere of Vandermond determinant.

4.5.1 Resolving $T_{2\alpha_2}$ discrete transformation chain

The $T_{2\alpha_2}$ discrete transformation may be presented as linear chain of equations for only two unknown functions $f^-_{1.1}, f^-_{1.2}$. Paramerizing them in the form $(f^-_{1.1})_n = \frac{A^n}{D^2_n}$, $\quad (f^-_{1.2})_n = \frac{B^n}{D^2_n}$ we pass to the following chain of equations

$$A^{n+1} = \frac{\frac{1}{2}D_{n+1}B^n_1 - B^n(D_{n+1})_1}{D_n} \tag{18}$$

$$\frac{B^{n+1}}{D^2_{n+1}} = \frac{1}{D^2_n}[\frac{1}{4}B^n_{11} - \frac{(D_{n+1})_1}{D_{n+1}}B^n_1 + (\frac{(D_{n+1})_1}{D_{n+1}})^2 B^n] + \tag{19}$$

$$\frac{A^n(D_{n+1})_1 - D_{n+1}A^n_1}{2D^3_n} + \frac{D_{n+1}(A^n(D_n)_1 + B^n D_{n-1})}{2D^4_n}$$

The last unknown function in these notations looks as $f^-_{1.0} = \frac{B^{n-1}}{D^2_n}$.

Let us resolve (19) by induction, assuming (see appendix), that

$$A^n = \frac{1}{n!(n+1)!}\int P(\lambda)d\lambda \prod_{i=1}^{(2n+1)} Q(\mu_i)d\mu_i \frac{W^2_n(\mu_1, ..\mu_n)W^2_{n+1}(\mu_{n+1}, ..\mu_{2n+1})}{\prod_{i=1}^{(2n+1)}(\lambda - \mu_i)} \tag{20}$$

$$B^n = \frac{1}{(n+1)!(n+1)!}\int P(\lambda)d\lambda \prod_{i=1}^{(2n+2)} Q(\mu_i)d\mu_i \frac{W^2_{n+1}(\mu_1, ..\mu_{n+1})W^2_{n+1}(\mu_{n+2}, ..\mu_{2n+2})}{\prod_{i=1}^{(2n+2)}(\lambda - \mu_i)} \tag{21}$$

From this moment all factorial factors are included in definition of Vandermond determinant. Substituting assumed form of solution into (18) we obtain integrant function in a form

$$\frac{W^2_{n+1}(\mu_1, ..\mu_{n+1})W^2_{n+1}(k_1, ..k_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda - \mu_i)(\lambda - k_i)}(\frac{1}{2}\sum_{i=1}^{n+1}(\mu_i + k_i) - \sum_{i=1}^{n+1} d_i)W^2_{n+1}(d_1, ...d_{n+1})$$

The sum with factor $\frac{1}{2}$ arises after differentiation B^n on $D_{1,0}$. The second sum arises after differentiation of D_{n+1} on the same operator. But domain of integration is symmetrical with respect $(2n+2)$ parameters (m, k) and $(n+1)$ parameters d. Thus it is possible rewrite the last expression in a form

$$\frac{W^2_{n+1}(\mu_1, ..\mu_{n+1})W^2_{n+1}(k_1, ..k_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda - \mu_i)(\lambda - k_i)}(\sum_1^{n+1}(\mu_i - d_i)W^2_{n+1}(d_1, ...d_{n+1}) \tag{22}$$

Now let us compare the last expression with solution in the case of A_2 arising after application n times discrete transformation T_2. For interesting for us functions in connection with (11) we have

$$f^-_{1.0} = \frac{V(1, n+1)}{V(0, n+1)} \quad , f^-_{1.1} = \frac{V(1, n+2)}{V(0, n+1)}, \quad f^+_{0.1} = \frac{V(0, n)}{V(0, n+1)}$$

We rewrite now equation for $f^-_{1.0}$ (6) conserving only integrant function in both sides

$$\frac{W^2_{n+1}(\mu_1,..\mu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\mu_i)}(\sum_{i=1}^{n+1}\mu_i-\sum_{i=1}^{n+1}d_i)W^2_{n+1}(d_1,....d_{n+1})=$$
$$\frac{W^2_{n+2}(\mu_1,..\mu_{n+2})}{\prod_{i=1}^{(n+2)}(\lambda-\mu_i)}W^2_n(d_1,....d_n)) \tag{23}$$

where $\mu_{n+2}=d_{n+1}$ (to have the same integrable indexes in both sides of the last equality). The last equality after substitution it into (22) finish the proof of reccurent relation for A^{n+1} term of the chain above.

Now we pass to calculation of B^{n+1} defined by (19). The terms in the second row of this equality may be rewrriten in a equivalent form

$$\frac{A^n(D_{n+1}D_n)_1-D_{n+1}D_nA^n_1+B^nD_{n+1}D_{n-1}}{2D^4_n}$$

Substituting into the last expression assuming by reduction form of A^n and using once more (23) we represent it as

$$-\frac{\int P(\lambda)d\lambda\prod_{i=1}^{(n+2)}Q(\mu_i)d\mu_i\prod_{i=1}^{n}Q(\nu_i)d\nu_i\frac{W^2_n(\nu_1,..\nu_n)W^2_{n+2}(\mu_1,..\mu_{n+2})}{\prod_{i=1}^{(n+2)}(\lambda-\mu_i)\prod_{k=1}^{n}(\lambda-\nu_k)}}{2D^2_n}$$

The summation the terms of the first row of (19) leads to the following result in numenator

$$\frac{W^2_{n+1}(\mu_1,..\mu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\mu_i)}\frac{W^2_{n+1}(\nu_1,....\nu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\mu_i)}$$
$$(\sum_1^{n+1}(\frac{\mu_i+\nu_i}{2}-\sigma_i)\sum_1^{n+1}(\frac{\mu_i+\nu_i}{2}-\delta_i)W^2_{n+1}(\sigma_1,..\sigma_{n+1})W^2_{n+1}(\delta_1,..\delta_{n+1}) \tag{24}$$

In denumerator we have $D^2_nD^2_{n+1}$. The origin of the terms in (24) is as follows. Each term contain linear B^n and its derivatives. We parametrised this function by ν,μ parameters. Each term contains second degree of D_{n+1} or its derivatives. We parametrised them by independent parameters σ,δ). The first derivatives of B^n function leads to multiplicator $\sum_1^{n+1}(\mu_i+\nu_i)$. Keeping in mind the factor $\frac{1}{4}$ at the first term in (19) we explain corresponding term in (24). Quadratical in derivades on $(D_{n+1})_1$ terms leads obviously to product of sums of σ and δ parameters and so on.

Now stratedgy of the further calculatios will be the following. We multiply both sides of (19) on D^2_n and use once more (23) in the back direction we come to the following expression in the left side:

$$\frac{W^2_{n+2}(\mu_1,..\mu_{n+2})}{\prod_{i=1}^{(n+2)}(\lambda-\mu_i)}W^2_n(\sigma_1,....\sigma_n)\frac{W^2_{n+2}(\nu_1,..\nu_{n+2})}{\prod_{i=1}^{(n+2)}(\lambda-\nu_i)}W^2_n(\delta_1,....\delta_n)$$

where determinant of Vandermond parametrized by μ, ν arised from the assumed form of B^{n+1} (21), parametrised by δ, σ ones from D_n^2. Now let us use (23) twise. We obtain

$$\frac{W^2_{n+1}(\mu_1,..\mu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\mu_i)}(\sum_{i=1}^{n+1}\mu_i-\sum_{i=1}^{n+1}\delta_i)W^2_{n+1}(\delta_1,....\delta_{n+1})\times$$

$$\frac{W^2_{n+1}(\nu_1,..\nu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\nu_i)}(\sum_{i=1}^{n+1}\nu_i-\sum_{i=1}^{n+1}\sigma_i)W^2_{n+1}(\sigma_1,....\sigma_{n+1})$$

The structure of last expression exactly the same as in (24)- the combination terms in the right side of (21). Different only multiplicators. Differenece of them is the following ($\mu \equiv \sum_{i=1}^{n+1}\mu_i$ and so on)

$$(\frac{\mu+\nu}{2}-\sigma)(\frac{\mu+\nu}{2}-\delta)-(\mu-\delta)(\nu-\sigma)=(\frac{\mu-\nu}{2})^2$$

In the last expression dependence on parameteres δ, σ is factorised and leads to D^2_{n+1} which is canceled with the same term in denumeratoor. The remaining relation, which have been proved is the following

$$\frac{W^2_{n+1}(\nu_1,..\nu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\nu_i)}(\frac{\sum_{i=1}^{n+1}\nu_i-\sum_{i=1}^{n+1}\mu_i}{2})^2\frac{W^2_{n+1}(\mu_1,....\mu_{n+1})}{\prod_{i=1}^{(n+1)}(\lambda-\mu_i)}-$$

$$-\frac{1}{2}\frac{W^2_n(\nu_1,..\nu_n)W^2_{n+2}(\mu_1,..\mu_{n+2})}{\prod_{i=1}^{(n+2)}(\lambda-\mu_i)\prod_{k=1}^{n}(\lambda-\nu_k)}=0$$

But the last equality exactly coincedes with equations of the Toda lattice (17) Thus the solution of the linear chain (18) and (19)is proved by induction. Now to this solution it is necessary applicate n_1 times discrete transformation of the first simple root T_1. We will not do corresponding calculations but in the next subsection will consider discrete transformation chain of the first simple root. From this consideration general form of solution will be obvious.

4.5.2 Resolving $T_{1.0}$ discrete transformation chain

As it was mentioned after explicit formulae of discrete transformation (13) after action of this transformation on the initial functions $f^+_{1.0}=f^+_{0.1}=f^+_{1.1}=f^+_{1.2}=0$ the last three functions conserved there zero values. The remaing equations describing arising chain looks as

$$f^+_{1.0}=\frac{1}{\tilde{f}^-_{1.0}},\quad \tilde{f}^-_{0.1}=\frac{\tilde{f}^-_{1.1}}{f^-_{1.0}},\quad \tilde{f}^-_{1.2}=f^-_{1.2}-\frac{(f^-_{1.1})^2}{\tilde{f}^-_{1.0}}$$

$$\tilde{f}^-_{1.1}=D_{1,2}f^-_{1.1}+f^-_{0.1}f^-_{1.0}-\frac{1}{2}f^-_{1.1}D_{1,2}\ln f^-_{1.0} \tag{25}$$

$$\frac{\tilde{f}^-_{1.0}}{f^-_{1.0}} = \frac{1}{4} D_{1,2} D_{1,2} \ln f^-_{1.0} + f^+_{1.0} f^-_{1.0}$$

Now it is necessary to take into acount that $D_{1,0} f^-_{1.0} = 0$ and in connection of equations of motion (12) $D_{1,1} f^-_{1.0} = -f^-_{0.1} f^-_{1.0}$ (the last term $f^-_{1.2} f^+_{0.1} = 0$) we rewrite the last two equations from (25) in a equivalent form

$$\tilde{f}^-_{1.1} = f^-_{1.0} D_{0,1} \frac{f^-_{1.1}}{f^-_{1.0}}$$

$$\frac{\tilde{f}^-_{1.0}}{f^-_{1.0}} = D^2_{0,1} \ln f^-_{1.0} + f^+_{1.0} f^-_{1.0}, \quad f^+_{1.0} = \frac{1}{\tilde{f}^-_{1.0}}$$

The system of equation above for $f^+_{1.0}, f^-_{1.0}, f^-_{1.1}$ functions exactly coinsides with discrete transformation T_1 (9) for the first simple root of A_2 algebra solition of which was found above in the third section. The last function $f^-_{1.2}$ is defined algebraicaly via $f^-_{1.1}, f^-_{1.0}$ from the first row of (25). By reduction using only first equality of Yacobi [2] we obtain

$$f^-_{1.1} = \frac{\tilde{D}et_{n+1}}{Det_n}, \quad f^-_{1.2} = \frac{\tilde{\tilde{D}}et_{n+1}}{Det_n} \tag{26}$$

where Det_n are the main minores of the matrix

$$\begin{pmatrix} f^-_{1.0} & D_{0,1} f^-_{1.0} & D^2_{0,1} f^-_{1.0} & ..... \\ D_{0,1} f^-_{1.0} & D^2_{0,1} f^-_{1.0} & D^3_{0,1} f^-_{1.0} & ..... \\ D^2_{0,1} f^-_{1.0} & D^3_{0,1} f^-_{1.0} & D^4_{0,1} f^-_{1.0} & ..... \\ ...... & .... & .... & ..... \end{pmatrix} \tag{27}$$

$\tilde{D}_n$ determinant of matrix (27) in which last column is constructed from $f^-_{1.1}$ function and its consequent derivatives. And at last $\tilde{\tilde{D}}_n$ in matrix (27) the last column and row are exchanged on consequent deritivies of $f^-_{1.1}$ function exept of diagonal term which is occupied by $f^-_{1.2}$ function. One example clarify the situation

$$\tilde{\tilde{D}}_3 = Det_3 \begin{pmatrix} f^-_{1.0} & D_{0,1} f^-_{1.0} & f^-_{1.1} \\ D_{0,1} f^-_{1.0} & D^2_{0,1} f^-_{1.0} & D_{0,1} f^-_{1.1} \\ f^-_{1.1} & D_{0,1} f^-_{1.1} & f^-_{1.2} \end{pmatrix}$$

Integrales in the (26) may be calculated directly by methods of [4]. We present calculations for more complicate case of $\tilde{\tilde{D}}et_{n+1}$. We paramerise first n elements of the first column by λ_1 (parameter of integration) and the last term $f^-_{1.1}$ by two parameters λ_1, μ_1. Elements of second column are parametrised by λ_2, μ_1 and so on up to $n - th$ column λ_n, μ_1. Elements of $(n+1)$-th column n first one by λ_{n+1}, μ_2 and the last one $f^-_{1.2}$ by three parameters $\lambda_{n+1}, \mu_1, \mu_2$. As a result under the sign of multidimensional $((n+3))$ integral we obtain determinant, which may be easely calculated by the same method as calculated Vandermond determinant. Result is as follows (we present integrand function)

$$\tilde{D}^{\tilde{}}_{n+1} = \frac{W^2_{n+1}(\lambda)}{\prod_{i=1}^{(n+1)}(\lambda_i-\mu_1)(\lambda_i-\mu_2)}, \quad \tilde{D}_{n+1} = \frac{W^2_{n+1}(\lambda)}{\prod_{i=1}^{(n+1)}(\lambda_i-\mu)}$$

Keeping in mind that transformation $T_{2\alpha_2}$ change only n_2 and transformation T_1 changes only n_1 we come to the solution of the next subsection

4.6 General formulae for solution

The result of calculation of discretre transformation $T^{n_2}_{2\alpha_2}T^{n_1}_1$ may be expessed in terms of the basis function

$$V(n_1;n_2,n_3) = \frac{1}{n_1!n_2!n_3!}\int\prod_{k=1}^{n_1}P(\lambda_k)d\lambda_k\prod_{i=1}^{n_2)}Q(\mu_i)d\mu_i\prod_{j=1}^{n_3)}Q(\nu_j)d\nu_j\times$$

$$\frac{W^2(\lambda)_{n_1}W^2(\mu)_{n_2}W^2(\nu)_{n_3}}{\prod_{k=1}^{n_1}\prod_{i=1}^{n_2}\prod_{j=1}^{n_3}(\lambda_k-\mu_i)(\lambda_k-\nu_j)}$$

In this notations solution of 4-wave B_2 problem looks similar as it was found before for the case of A_2 algebra

$$\begin{gathered}
f^+_{1.0} = \frac{V(n_1-1;n_2,n_2)}{V(n_1;n_2,n_2)}, \quad f^+_{0.1} = \frac{V(n_1;n_2,n_2-1)}{V(n_1;n_2,n_2)} \\
f^+_{1.1} = \frac{V(n_1-1;n_2,n_2-1)}{V(n_1;n_2,n_2)}, \quad f^+_{1.2} = \frac{V(n_1-1;n_2-1,n_2-1)}{V(n_1;n_2,n_2)} \\
f^-_{1.0} = \frac{V(n_1+1;n_2,n_2)}{V(n_1;n_2,n_2)}, \quad f^-_{0.1} = \frac{V(n_1;n_2+1,n_2)}{V(n_1;n_2,n_2)} \\
f^-_{1.1} = \frac{V(n_1+1;n_2+1,n_2)}{V(n_1;n_2,n_2)}, \quad f^-_{1.2} = \frac{V(n_1+1;n_2+1,n_2+1)}{V(n_1;n_2,n_2)}
\end{gathered} \tag{28}$$

In the case $n_1=0$ this exactly solution of chain of the second root; in the case $n_2=0$ this is solution of the chain of the first root.

5 The case of G_2 algebra

Algebra G_2 has the following Cartan matrix and basic commutation relations between two generators of the simple roots $X^{\pm}_{1,2}$ and its Cartan elements $h_{1,2}$

$$k = \begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}, \quad \begin{pmatrix} [h_1,X^{\pm}_1] = \pm 2X^{\pm}_1 & [h_1,X^{\pm}_2] = \mp X^{\pm}_2 \\ [h_2,X^{\pm}_1] = \mp 3X^{\pm}_2 & [h_2,X^{\pm}_2] = \pm 2X^{\pm}_2 \end{pmatrix}$$

System of its positive roots contain 6 elements

$$X_1^+ = X_{\alpha_1}^+, X_2^+ = X_{\alpha_2}^+, X_{1;2}^+ = [X_2^+, X_1^+] = X_{\alpha_1+\alpha_2}^+, X_{1;22}^+ = \frac{1}{2}[X_2^+[X_2^+, X_1^+]] = X_{\alpha_1+2\alpha_2}^+,$$

$$X_{1;222}^+ == \frac{1}{6}[X_2^+[X_2^+[X_2^+, X_1^+]]] = X_{\alpha_1+3\alpha_2}^+, X_{11;222}^+ = \frac{1}{6}[X_1^+[X_2^+[X_2^+[X_2^+, X_1^+]]]] = X_{2\alpha_1+3\alpha_2}^+$$

The coefficients of decomposition by these roots will be denoted as $f_{n;m}$ correspondingly. System of equations for 12 functions $f_{2.3}^\pm, f_{1.3}^\pm, f_{1.2}^\pm, f_{1.1}^\pm, f_{1.0}^\pm, f_{0.1}^\pm$ looks as

$$\begin{aligned}
D_{2,3}f_{2.3}^+ &= 3f_{1.0}^+f_{1.3}^+ - 3f_{1.1}^+f_{1.2}^+, \quad D_{2,3}f_{2.3}^- = 3f_{1.0}^-f_{1.3}^- - 3f_{1.1}^-f_{1.2}^- \\
D_{1,3}f_{1.3}^+ &= -3f_{2.3}^+f_{1.0}^- - 3f_{0.1}^+f_{1.2}^+, \quad D_{1,3}f_{1.3}^- = -3f_{1.0}^+f_{2.3}^- - 3f_{0.1}^-f_{1.2}^- \\
D_{1,2}f_{1.2}^+ &= f_{2.3}^+f_{1.1}^- + f_{1.3}^+f_{0.1}^- - 2f_{0.1}^+f_{1.1}^+, \quad D_{1,2}f_{1.2}^- = f_{2.3}^-f_{1.1}^+ + f_{1.3}^-f_{0.1}^+ - 2f_{0.1}^-f_{1.1}^- \\
&\qquad (29) \\
D_{1,1}f_{1.1}^+ &= f_{2.3}^+f_{1.2}^- + 2f_{1.2}^+f_{0.1}^- - f_{0.1}^+f_{1.0}^+, \quad D_{1,1}f_{1.1}^- = f_{2.3}^-f_{1.2}^+ + 2f_{1.2}^-f_{0.1}^+ - f_{0.1}^-f_{1.0}^- \\
D_{1,0}f_{1.0}^+ &= -3f_{2.3}^+f_{1.3}^- + 3f_{1.1}^+f_{0.1}^-, \quad D_{1,0}f_{1.0}^- = -3f_{2.3}^-f_{1.3}^+ + 3f_{1.1}^-f_{0.1}^+ \\
D_{0,1}f_{0.1}^+ &= f_{1.3}^+f_{1.2}^- + 2f_{1.2}^+f_{1.1}^- + f_{1.1}^+f_{1.0}^-, \quad D_{0,1}f_{0.1}^- = f_{1.3}^-f_{1.2}^+ + 2f_{1.2}^-f_{1.1}^+ + f_{1.1}^-f_{1.0}^+
\end{aligned}$$

5.1 Initial conditions for G_2

Let us rewrite (3) and (4) in a form under condition that all $f^+ = 0$

$$f_{1.0}^- = \int d\lambda p(\lambda)e^{\lambda(c_1x-d_1t)}, \quad f_{0.1}^- = \int d\mu q(\mu)e^{\mu(c_2x-d_2t)},$$

$$f_{1.1}^- = \int d\lambda \int d\mu \frac{p(\lambda)q(\mu)}{\lambda-\mu} e^{\lambda(c_1x-d_1t)+\mu(c_2x-d_2t)}$$

$$f_{1.2}^- = -\int d\lambda \int d\mu_1 d\mu_2 \frac{p(\lambda)q(\mu_1)q(\mu_2)}{(\lambda-\mu_1)(\lambda-\mu_2)} e^{\lambda(c_1x-d_1t)+(\mu_1+\mu_2)(c_2x-d_2t)}$$

$$f_{1.3}^- = \int d\lambda \int d\mu_1 d\mu_2 d\mu_3 \frac{p(\lambda)q(\mu_1)q(\mu_2)q(\mu_3)}{(\lambda-\mu_1)(\lambda-\mu_2)(\lambda-\mu_3)} e^{\lambda(c_1x-d_1t)+(\mu_1+\mu_2+\mu_3)(c_2x-d_2t)}$$

$$f_{2.3}^- = \frac{1}{2}\int d\lambda_1 d\lambda_2 (\lambda_1-\lambda_2)^2 \int d\mu_1 d\mu_2 d\mu_3 \frac{p(\lambda_1)p(\lambda_2)q(\mu_1)q(\mu_2)q(\mu_3)}{\prod_{i=1,a=1}^{i=2,a=3}(\lambda_i-\mu_a)} e^R$$

$$R = (\lambda_1+\lambda_2)(c_1x-d_1t) + (\mu_1+\mu_2+\mu_3)(c_2x-d_2t)$$

5.2 Discrete transformation of the first simple root T_1

We remind to the reader that first seven equation of discrete transformation below is a direct consequence of results of the paper [1]. Of course it is possible consider them as a happy guess. The most simple way of obtaining last five equation consists in direct substitution the 7 previous ones in the system (29).

$$\tilde{f}^+_{1.0} = \frac{1}{f^-_{1.0}}, \quad \tilde{f}^-_{1.3} = -\frac{f^-_{2.3}}{f^-_{1.0}}, \quad \tilde{f}^+_{1.1} = -\frac{f^+_{0.1}}{f^-_{1.0}}, \quad \tilde{f}^-_{0.1} = \frac{f^-_{1.1}}{f^-_{1.0}}, \quad \tilde{f}^+_{2.3} = \frac{f^+_{1.3}}{f^-_{1.0}}$$

$$\tilde{f}^+_{1.2} = f^+_{1.2} + \frac{f^-_{1.1}f^+_{1.3} + (f^+_{0.1})^2}{f^-_{1.0}}, \quad \tilde{f}^-_{1.2} = f^-_{1.2} - \frac{f^-_{2.3}f^+_{0.1} + (f^-_{1.1})^2}{f^-_{1.0}}$$

The expressions above now it is possible to substitute into (29) and obtain all other equations of discrete transformation. For instant rewriting equation for $f^-_{1.3}$ function for transformed variables

$$D_{1,3}\tilde{f}^-_{1.3} = -3\tilde{f}^+_{1.0}\tilde{f}^-_{2.3} - 3\tilde{f}^-_{0.1}\tilde{f}^-_{1.2}$$

and substituting in in it transformed functions from above equations after not combersome calculations we find $\tilde{f}^-_{2.3}$ and so on with the result

$$\tilde{f}^-_{2.3} = \frac{f^-_{1.0}D_{1,2}f^-_{2.3} - \frac{1}{2}f^-_{2.3}D_{1,2}f^-_{1.0}}{f^-_{1.0}} - f^-_{1.0}f^-_{1.3} + \frac{-(f^-_{2.3})^2 f^+_{1.3} + 2(f^-_{1.1})^3 + 3f^-_{2.3}f^-_{1.1}f^+_{0.1}}{2f^-_{1.0}}$$

$$\tilde{f}^+_{0.1} = \frac{f^-_{1.0}D_{1,2}f^+_{0.1} - \frac{1}{2}f^+_{0.1}D_{1,2}f^-_{1.0}}{f^-_{1.0}} - f^+_{1.1}f^-_{1.0} + \frac{2(f^-_{1.1})^2 f^+_{1.3} + (f^+_{0.1})^2 f^-_{1.1} + f^-_{2.3}f^+_{0.1}f^+_{1.3}}{2f^-_{1.0}}$$

$$\tilde{f}^+_{1.3} = \frac{f^-_{1.0}D_{1,2}f^+_{1.3} - \frac{1}{2}f^+_{1.3}D_{1,2}f^-_{1.0})}{f^-_{1.0}} + f^+_{2.3}f^-_{1.0} + \frac{(f^+_{1.3})^2 f^-_{2.3} - 3f^+_{1.3}f^-_{1.1}f^+_{0.1} - (f^+_{0.1})^3}{2f^-_{1.0}}$$

$$\tilde{f}^-_{1.1} = \frac{f^-_{1.0}D_{1,2}f^-_{1.1} - \frac{1}{2}f^-_{1.1}D_{1,2}f^-_{1.0}}{f^-_{1.0}} + f^-_{0.1}f^-_{1.0} - \frac{2(f^+_{0.1})^2 f^-_{2.3} + (f^-_{1.1})^2 f^+_{0.1} + f^-_{2.3}f^-_{1.1}f^+_{1.3}}{2f^-_{1.0}}$$

$$\frac{\tilde{f}^-_{1.0}}{f^-_{1.0}} = \frac{1}{4}D^2_{1,2}\ln f^-_{1.0} + f^-_{1.0}f^+_{1.0} + \frac{1}{2}(f^-_{0.1}f^+_{0.1} + f^-_{1.1}f^+_{1.1}) + \frac{3}{2}(f^-_{2.3}f^+_{2.3} + f^-_{1.3}f^+_{1.3}) +$$

$$\frac{3}{4}\frac{f^+_{0.1}D_{1,2}f^-_{1.1} - f^-_{1.1}D_{1,2}f^+_{0.1}}{f^-_{1.0}} - \frac{1}{4}\frac{f^+_{1.3}D_{1,2}f^-_{2.3} - f^-_{2.3}D_{1,2}f^+_{1.3}}{f^-_{1.0}} - \frac{f^+_{0.1}}{f^-_{1.0}}(f^-_{2.3}f^+_{1.2} - f^+_{0.1}f^-_{1.2}) -$$

$$\frac{1}{4}\frac{3(f^-_{1.0}f^+_{0.1})^2 - (f^-_{2.3}f^+_{1.3})^2 + 6f^-_{1.0}f^+_{0.1})f^-_{2.3}f^+_{1.3} + 4(f^-_{1.1})^3 f^+_{1.3} + 4(f^+_{0.1})^3 f^-_{2.3}}{(f^-_{1.0})^2}$$

5.3 Discrete transformation of the complicate root $T_{\alpha_1+3\alpha_2}$

The calculations below are not necessary if one pay attention on symmetry of the main system of equations (29) with respect to the following exchange of variables and unknown functions

$$D_{2,3} \to -D_{2,3}, \quad D_{1.3} \to -D_{1,0}, \quad D_{1,2} \to -D_{1,1}, \quad D_{0,1} \to D_{0,1},$$

$$f^{\pm}_{2.3} \to -f^{\pm}_{2.3}, \quad f^{\pm}_{1.3} \to -f^{\pm}_{1.0}, \quad f^{\pm}_{1.1} \to -f^{\pm}_{1.2}, \quad f^{\pm}_{0.1} \to -f^{\mp}_{0.1}$$

Using this substitution or by straitforward calculations for $T_{\alpha_1+3\alpha_2}$ we obtain

$$\tilde{f}^{+}_{1.3} = \frac{1}{f^{-}_{1.3}}, \quad \tilde{f}^{+}_{0.1} = \frac{f^{-}_{1.2}}{f^{-}_{1.3}}, \quad \tilde{f}^{+}_{1.2} = -\frac{f^{-}_{0.1}}{f^{-}_{1.3}}, \quad \tilde{f}^{-}_{1.0} = \frac{f^{-}_{2.3}}{f^{-}_{1.3}}, \quad \tilde{f}^{+}_{2.3} = -\frac{f^{+}_{1.0}}{f^{-}_{1.3}}$$

$$\tilde{f}^{+}_{1.1} = f^{+}_{1.1} + \frac{(f^{-}_{0.1})^2 + f^{-}_{1.2}f^{+}_{1.0}}{f^{-}_{1.3}}, \quad \tilde{f}^{-}_{1.1} = f^{-}_{1.1} + \frac{-(f^{-}_{1.2})^2 + f^{-}_{2.3}f^{-}_{0.1}}{f^{-}_{1.3}}$$

$$\tilde{f}^{-}_{2.3} = \frac{-f^{-}_{1.3}D_{1,1}f^{-}_{2.3} + \frac{1}{2}f^{-}_{2.3}D_{1,1}f^{-}_{1.3}}{f^{-}_{1.3}} + f^{-}_{1.0}f^{-}_{1.3} + \frac{(f^{-}_{2.3})^2 f^{+}_{1.0} - (f^{-}_{1.2})^3 + 3f^{-}_{2.3}f^{-}_{1.1}f^{-}_{1.2}}{2f^{-}_{1.3}}$$

$$\tilde{f}^{+}_{1.0} = \frac{-f^{-}_{1.3}D_{1,1}f^{+}_{1.0} + \frac{1}{2}f^{+}_{1.0}D_{1,1}f^{-}_{1.3}}{f^{-}_{1.3}} - f^{+}_{2.3}f^{-}_{1.3} - \frac{f^{-}_{2.3}(f^{+}_{1.0})^2 + 2(f^{-}_{0.1})^3 + 3f^{-}_{0.1}f^{-}_{1.2}f^{+}_{1.2}}{2f^{-}_{1.3}}$$

$$\tilde{f}^{-}_{1.2} = \frac{-f^{-}_{1.3}D_{1,1}f^{-}_{1.2} + \frac{1}{2}f^{-}_{1.2}D_{1,1}f^{-}_{1.3}}{f^{-}_{1.3}} - f^{+}_{0.1}f^{-}_{1.3} + \frac{f^{-}_{2.3}(f^{-}_{0.1})^2 + 2(f^{-}_{0.1})^2 f^{-}_{2.3} - f^{-}_{0.1}(f^{-}_{1.2})^2}{2f^{-}_{1.3}}$$

$$\tilde{f}^{-}_{0.1} = \frac{-f^{-}_{1.3}D_{1,1}f^{-}_{0.1} + \frac{1}{2}f^{-}_{0.1}D_{1,1}f^{-}_{1.3}}{f^{-}_{1.3}} - f^{+}_{1.2}f^{-}_{1.3} + \frac{f^{-}_{1.2}(f^{-}_{0.1})^2 + 2(f^{-}_{1.2})^2 f^{+}_{1.0} - f^{-}_{0.1}f^{-}_{2.3}f^{+}_{1.0})}{2f^{-}_{1.3}}$$

$$\frac{\tilde{f}^{-}_{1.3}}{f^{-}_{1.3}} = \frac{1}{4}D^2_{1,1}\ln f^{-}_{1.3} + f^{-}_{1.1}f^{+}_{1.3} + \frac{1}{2}(f^{-}_{0.1}f^{+}_{0.1} + f^{-}_{1.2}f^{+}_{1.2}) + \frac{3}{2}(f^{-}_{2.3}f^{+}_{2.3} + f^{-}_{1.0}f^{+}_{1.0}) -$$

$$\frac{3}{4}\frac{f^{-}_{0.1}D_{1,1}f^{-}_{1.2} - f^{-}_{1.2}D_{1,1}f^{-}_{0.1}}{f^{-}_{1.3}} - \frac{1}{4}\frac{f^{+}_{1.0}D_{1,1}f^{-}_{2.3} - f^{-}_{2.3}D_{1,1}f^{+}_{1.0}}{f^{-}_{1.3}} + \frac{f^{-}_{0.1}}{f^{-}_{1.3}}(f^{-}_{2.3}f^{+}_{1.1} + f^{-}_{0.1}f^{-}_{1.1}) -$$

$$\frac{1}{4}\frac{3(f^{-}_{1.3}f^{-}_{0.1})^2 - (f^{-}_{2.3}f^{+}_{1.0})^2 - 6f^{-}_{1.3}f^{-}_{0.1})f^{-}_{2.3}f^{+}_{1.0} + 4(f^{-}_{1.2})^3 f^{+}_{1.3} - 4(f^{-}_{0.1})^3 f^{-}_{2.3}}{(f^{-}_{1.3})^2}$$

Of course all formulae above for T_1 and $T_{\alpha_1+3\alpha_2}$ coinsides with the general once in the case of arbitrary semisimple algebra of the paper [1].

5.4 General formulae of solution

In spite of comlicate on the first look structure of dicrete transformations of the previous two subsections their resolving is observable. We would not like in this paper give the proofs but present only finally result of not simple calculations. Aritrary discrete transformation can be represented as consequent applications of two mutualy commutative basis transformations T_1 and $T_{3\alpha_2} \equiv T_1^{-1} T_{\alpha_1+3\alpha_2}$ and has the form $T = T_1^{n_1} T_{3\alpha_2}^{n_2}$. The result of calculation of discretre transformation $T_{3\alpha_2}^{n_2} T_1^{n_1}$ may be expessed in terms of the basis function (all factorial factors are included in definition of Wandermond determinants)

$$V(n_1; n_2, n_3, n_4) = \int \prod_{k=1}^{n_1} P(\lambda_k) d\lambda_k \prod_{i=1}^{n_2} Q(\mu_i) d\mu_i \prod_{j=1}^{n_3} Q(\nu_j) d\nu_j \prod_{l=1}^{n_4} Q(\sigma_l) d\sigma_l \times$$

$$\frac{W^2(\lambda)_{n_1} W^2(\mu)_{n_2} W^2(\nu)_{n_3} W^2(\sigma)_{n_4}}{\prod_{k=1}^{n_1} \prod_{i=1}^{n_2} \prod_{j=1}^{n_3} \prod_{l=1}^{n_4} (\lambda_k - \mu_i)(\lambda_k - \nu_j)(\lambda_k - \sigma_l)}$$

In this notations solution of 6-wave G_2 problem looks similar as it was found before for the case of A_2 and B_2 algebras

$$\begin{gathered}
f^+_{1.0} = \frac{V(n_1-1; n_2, n_2, n_2)}{V(n_1; n_2, n_2, n_2)}, \quad f^+_{0.1} = \frac{V(n_1; n_2-1, n_2, n_2)}{V(n_1; n_2, n_2, n_2)} \\
f^+_{1.1} = \frac{V(n_1-1; n_2-1, n_2, n_2)}{V(n_1; n_2, n_2, n_2)}, \quad f^+_{1.2} = \frac{V(n_1-1; n_2-1, n_2-1, n_2)}{V(n_1; n_2, n_2, n_2)} \\
f^+_{1.3} = \frac{V(n_1-1; n_2-1, n_2-1, n_2-1)}{V(n_1; n_2, n_2, n_2)}, \quad f^+_{2.3} = \frac{V(n_1-2; n_2-1, n_2-1, n_2-1)}{V(n_1; n_2, n_2, n_2)} \\
f^-_{1.0} = \frac{V(n_1+1; n_2, n_2, n_2)}{V(n_1; n_2, n_2, n_2)}, \quad f^-_{0.1} = \frac{V(n_1; n_2+1, n_2, n_2)}{V(n_1; n_2, n_2, n_2)} \\
f^-_{1.1} = \frac{V(n_1+1; n_2+1, n_2, n_2)}{V(n_1; n_2, n_2, n_2)}, \quad f^-_{1.2} = \frac{V(n_1+1; n_2+1, n_2+1, n_2)}{V(n_1; n_2, n_2, n_2)} \\
f^-_{1.2} = \frac{V(n_1+1; n_2+1, n_2+1, n_2)}{V(n_1; n_2, n_2, n_2)}, \quad f^-_{1.3} = \frac{V(n_1+1; n_2+1, n_2+1, n_2+1)}{V(n_1; n_2, n_2, n_2)} \\
f^-_{2.3} = \frac{V(n_1+2; n_2+1, n_2+1, n_2+1)}{V(n_1; n_2, n_2, n_2)}
\end{gathered} \tag{30}$$

Of course the formulae above on the level of the present paper it isnecessary consider as hipothesis and possible happy guess for solution of problem in the case of arbitrary semisimple algebra.

6 Multisoliton solutions

The explicit form of solution in the case of B_2 and G_2 allow to find conditions of the chain interupting and construct multisoliton solutions as it was done in the case of A_2 algebra [3],[4]

7 Outlook

The main result of the present paper consits in assuarence that calculations of this kind may done in the case of arbitrary semisimple algebra. It is possible to assume that to this aim is sufficient to resolve the discrete transformation chains of the simple roots of the algebra. The finally result will be obtained by simple multiplications with taking into account only differences of spectral parameters of simple roots connected on the sheam of Dynkin.

The most intrigued is the fact that soliton solutions of multidimensional integrable system leads to the same limitations on paramaters amplituda-phase (with correction on interaction) as in the case A_1 algebra.

The same surprising is existence of universal integrable chains in the field of Toda lattice. Of course there description and understanding of their nature is very interesting problem stated by the present paper.

Ussualy all results in the theory of integrable system may be interpreted as some equalities form the theory of representtion of semisimple algebras (relations between highest vector of different irreducible representations and so on). What kind problems of representation theory (if any) may be connected with arised equalities of the present paper?

At last the parameters λ, μ, ν_i (the last after taking into account homogineous part solutions (section 2) of all elements of lower triangular algebra) in comparisen with A_1 algebra case play role of spectral parameters. By this reason it is possible to assume that one-dimentional Lax formalism it is necessary to change on multidimensional one. This assumtion it is possible to put into connection with the fact that discrete transformation are always commutative and number of basis transformation coincides with the rank of the algebra.

All this problems demand for its solution further investigation.

8 Appendix

In this Appendix we present detail calculations for the most complicate case of discrete transformation $T_{2\alpha_2}$ of B_2 algebra. We change greek indexes on latina ones here.

8.1 Initial condition. Zero order case

$$A^0 = \int \frac{dmMdlL}{l-m}, \quad B^0 = \int \frac{dm_1M_1dm_2M_2dlL}{(l-m_1)(l-m_2)}$$

8.2 First step of $T_{2\alpha_2}$ transformation

$$A^1 = \frac{\frac{1}{2}D_1B^0_1 - B^0(D_1)_1}{D_0} = \int \frac{dm_1M_1dm_2M_2dm_3M_3dlL(\frac{m_1+m_2}{2} - m_3)}{(l-m_1)(l-m_2)} =$$

$$\int\frac{dm_1M_1dm_2M_2dm_3M_3dlL(m_1-m_3)}{(l-m_1)(l-m_2)}=$$

$$\frac{1}{2}\int dm_1M_1dm_2M_2dm_3M_3dlL\frac{(m_1-m_3)}{(l-m_2)}(\frac{1}{l-m_1}-\frac{1}{l-m_3})=$$

$$\frac{1}{2}\int dm_1M_1dm_2M_2dm_3M_3dlL\frac{(m_1-m_3)^2}{(l-m_2)(l-m_1)(l-m_3)}$$

In transformation above it was used only the fact the symmetry of the domain of integration with respect to permutation of all indexes m_1,m_2,m_3. Under calculation of B^1 from (19) we will not conserve indexes of integrale and differentiales remaining only integrant function. We have consequently

$$\frac{1}{4}\frac{(m_1+m_2)^2}{(l-m_1)(l-m_2)}+\frac{m_4(m_3-(m_1+m_2))}{(l-m_1)(l-m_2)}-\frac{1}{2}\frac{(m_1-m_2)}{(l-m_1)}=$$

$$\frac{m_1(m_2-m_4)+m_4(m_3-m_2)}{(l-m_1)(l-m_2)}=\frac{1}{2}[\frac{m_1(m_2-m_4)^2}{(l-m_1)(l-m_2)(l-m_4)}-\frac{m_4(m_2-m_3)^2}{(l-m_1)(l-m_2)(l-m_3)}=$$

$$\frac{1}{2}\frac{(m_1-m_4)(m_2-m_3)^2}{(l-m_1)(l-m_2)(l-m_3)}=\frac{1}{4}\frac{(m_1-m_4)^2(m_2-m_3)^2}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)}$$

And thus for B^1 we obtain

$$B^1=\frac{1}{4}\int dm_1M_1dm_2M_2dm_3M_3dm_4M_4dlL\frac{(m_1-m_4)^2(m_2-m_3)^2}{(l-m_2)(l-m_1)(l-m_3)(l-m_4)}$$

In the process of calculations above we have used only the fact symmetry of region of integration with respect to permutations of four parameters m_i

$$A^2=\frac{1}{8}\frac{(m_5-m_6)^2(m_1-m_4)^2(m_2-m_3)^2}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)}[\frac{1}{2}(m_1+m_2+m_3+m_4)-(m_5+m_6)]=$$

$$\frac{1}{4}\frac{(m_5-m_6)^2(m_1-m_4)^2(m_2-m_3)^2}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)}[m_1-m_5]$$

Now it is necessary to use equality

$$(m_i-m_k)(m_j-m_l)=(m_j-m_k)(m_i-m_l)-(m_j-m_i)(m_k-m_l)$$

with help of which we come to the equality

$$\frac{1}{4}\frac{(m_5-m_6)^2(m_1-m_4)^2(m_2-m_3)^2}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)}[m_1-m_5]=$$

$$\frac{1}{2}\frac{(m_2-m_3)^2(m_1-m_4)(m_5-m_6)(m_1-m_6)(m_5-m_4)(m_1-m_5)}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)}=$$

$$\frac{1}{2}\frac{(m_2-m_3)^2(m_5-m_6)(m_1-m_6)(l-m_5)W_3(m_1,m_4,m_5)}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)(l-m_5)}=$$

$$\frac{1}{2!3!}\frac{(m_2-m_3)^2W_3^2(m_1,m_4,m_5)}{(l-m_1)(l-m_2)(l-m_3)(l-m_4)(l-m_5)}$$

By induction we obtain

$$A^n=\frac{1}{n!(n+1)!}\int Ldl\prod_{i=1}^{(2n+1)}M_idm_i\frac{W_n^2(m_1,..m_n)W_{n+1}^2(m_{n+1},..m_{2n+1})}{\prod_{i=1}^{(2n+1)}(l-m_i)}$$

$$B^n=\frac{1}{(n+1)!(n+1)!}\int Ldl\prod_{i=1}^{(2n+2)}M_idm_i\frac{W_{n+1}^2(m_1,..m_{n+1})W_{n+1}^2(m_{n+2},..m_{2n+2})}{\prod_{i=1}^{(2n+2)}(l-m_i)}$$

References

[1] A.N.Leznov *Theor. Math. Physica N132 (1): (955-969), 2002.*

[2] A.N.Leznov and R.Torres-Cordoba *Journal of Math. Phys. N123,v 2, (633-650), MAY 2000.*

[3] A.N.Leznov G.R.Toker and R.Torres-Cordoba *hep-th/0605009*

[4] A.N.Leznov G.R.Toker and R.Torres-Cordoba *Nonlinear Mathematical Physics, to be published*

[5] A.N.Leznov *Theor. Math. Physica N123,v 2, (633-650), MAY 2000.*

[6] A.N.Leznov *Physics of elementary particals and atom nuclears N27, v.5, p 1161-1246.*

[7] A.N.Leznov and M.V.Saveliev *Group-theoretical methods for integration of nonlinear dynamical systems* Progress in Physics 15, Basel, (1992)

[8] A.N.Leznov and E.A.Yusbashjan *LMP v35, p. 345-349, (1995)*